

EFFECTIVE GENERICITY AND DIFFERENTIABILITY

RUTGER KUYPER AND SEBASTIAAN A. TERWIJN

ABSTRACT. We prove that a real x is 1-generic if and only if every differentiable computable function has continuous derivative at x . This provides a counterpart to recent results connecting effective notions of randomness with differentiability. We also consider multiply differentiable computable functions and polynomial time computable functions.

1. INTRODUCTION

The notion of 1-genericity is an effective notion of genericity from computability theory that has been studied extensively, see e.g. Jockusch [6], or the textbooks Odifreddi [11] and Downey and Hirschfeldt [5]. 1-Genericity, or Σ_1^0 -genericity in full, can be defined using computably enumerable (c.e.) sets of strings as forcing conditions. This notion captures a certain type of effective finite extension constructions that is common in computability theory. In this paper we give an characterization of 1-genericity in terms of familiar notions from computable analysis. This complements recent results by Brattka, Miller, and Nies [1] that characterize various notions of algorithmic randomness in terms of computable analysis. For example, in [1] it was proven (building on earlier work by Demuth [4]) that an element $x \in [0, 1]$ is Martin-Löf random if and only if every computable function of bounded variation is differentiable at x . Note that the notion of Martin-Löf randomness, which one could also call Σ_1^0 -randomness, is the measure-theoretic counterpart of the topological notion of 1-genericity.

The main result of this paper is as follows.

Theorem 1.1. *A real $x \in [0, 1]$ is 1-generic if and only if every differentiable computable function $f : [0, 1] \rightarrow \mathbb{R}$ has continuous derivative at x .*

The two implications of this theorem will be proven in Theorems 4.3 and 5.2. Note that by “differentiable computable function” we mean a computable function that is classically differentiable, so that in particular the derivative need not be continuous. Our result can be seen an effectivization of a result by Bruckner and Leonard.

Theorem 1.2. (Bruckner and Leonard [3, p. 27]) *A set $A \subseteq \mathbb{R}$ is the set of discontinuities of a derivative if and only if A is a meager Σ_2^0 set.*

One might expect that, in analogy to Theorem 1.1, n times differentiable computable functions would characterize n -genericity. However, in section 7 we show that 1-genericity is also equivalent to the n th derivative of any n times differentiable computable function being continuous at x . In section 8 we consider differentiable polynomial time computable functions and show that again these characterize 1-genericity.

Date: August 26, 2014.

The research of the first author was supported by NWO/DIAMANT grant 613.009.011 and by John Templeton Foundation grant 15619: ‘Mind, Mechanism and Mathematics: Turing Centenary Research Project’.

Our notation is mostly standard. We denote the natural numbers by ω . The Cantor space of all infinite binary sequences is denoted by 2^ω , and $2^{<\omega}$ is the set of all finite binary strings. For a finite string σ and a finite or infinite string x , we denote by $\sigma \sqsubseteq x$ that σ is an initial segment of x . For a string $\sigma \in 2^{<\omega}$, we have

$$[\sigma] = \{x \in 2^\omega : \sigma \sqsubseteq x\}.$$

The product topology on 2^ω , sometimes called the tree topology, or the finite information topology, has all sets of the form $[\sigma]$ as basic open sets. For a set $A \subseteq 2^{<\omega}$, we let

$$[A] = \bigcup_{\sigma \in A} [\sigma].$$

Thus every set A of finite strings defines an open subset of 2^ω . A subset of 2^ω is a Σ_1^0 class, or *effectively open*, if it is of the form $[A]$, with $A \subseteq 2^{<\omega}$ computably enumerable (c.e.). A set is a Π_1^0 class, or *effectively closed*, if it is the complement of a Σ_1^0 class. Thus Σ_1^0 and Π_1^0 classes form the first levels of the effective Borel hierarchy. As usual, the levels of the classical Borel hierarchy are denoted by boldface symbols Σ_n^0 and Π_n^0 . These notions are defined in the same way for $[0, 1]$, using rational intervals as basic opens. We denote the interior of a set $V \subseteq 2^\omega$ by $\text{Int}(V)$.

For unexplained notions from computability theory, we refer to Odifreddi [10] or Downey and Hirschfeldt [5]. For background in descriptive set theory we refer to Kechris [7] or Moschovakis [9]. Further background on (classical) Baire category theory can also be found in Oxtoby [12].

2. 1-GENERICITY

First, let us recall what it means for an element $x \in 2^\omega$ to be 1-generic. We will then discuss 1-genericity for elements of $[0, 1]$. A discussion of the properties of arithmetically generic and 1-generic sets can be found in Jockusch [6]. The “forcing-free” formulation of genericity we use here is due to Posner, see [6, p115].

Given a sequence $x \in 2^\omega$ and a set $A \subseteq 2^{<\omega}$, we say that x *meets* A if there exists $\sigma \sqsubseteq x$ such that $\sigma \in A$; equivalently, if $x \in [A]$. The set A is *dense along* x if for every $\sigma \sqsubseteq x$ there is an extension $\tau \sqsupseteq \sigma$ such that $[\tau] \subseteq [A]$; equivalently, if x is in the closure of the open set $[A]$.

Definition 2.1. An element $x \in 2^\omega$ is *1-generic* if x meets every c.e. set $A \subseteq 2^{<\omega}$ that is dense along x .

We now reformulate the definition of 1-genericity into a form that will be convenient in what follows. This formulation is also better suited for the discussion of generic real numbers (as opposed to infinite strings).

Lemma 2.2. *Let $A \subseteq 2^{<\omega}$ and let $V = 2^\omega \setminus [A]$. Then A is dense along x if and only if x is not in the interior of V . Therefore, A is dense along x and x does not meet A if and only if $x \in V \setminus \text{Int}(V)$.*

Proof. Note that A is dense along x if and only if every open set containing x has nonempty intersection with $[A]$. Thus, A is dense along x if and only if every open set disjoint from $[A]$ does not contain x . However, the open sets disjoint from $[A]$ are exactly the open sets contained in V , of which $\text{Int}(V)$ is the largest. Thus A is dense along x if and only if $\text{Int}(V)$ does not contain x . \square

Corollary 2.3. *For any $x \in 2^\omega$ we have that x is 1-generic if and only if for every Π_1^0 class $V \subseteq 2^\omega$ we have $x \notin V \setminus \text{Int}(V)$.*

One of the reasons this is interesting to mention explicitly is because a typical example of a nowhere dense set is a closed set with its interior removed, and the Π_1^0 sets are the simplest type of closed sets. Thus, the Corollary 2.3 says that x

is 1-generic if it is not in any of the simple, typical nowhere dense sets. This way of looking at 1-generic sets complements the usual motivation of 1-genericity by forcing, and it also allows one to easily compare 1-genericity with weak 1-genericity (since x is weakly 1-generic if it is not in any Π_1^0 -class with empty interior, see [5]).

With this equivalence in mind, we can now also define what it means for an element of $[0, 1]$ to be 1-generic.

Definition 2.4. Let $x \in [0, 1]$. We say that x is 1-generic if for every Π_1^0 class $V \subseteq [0, 1]$ we have $x \notin V \setminus \text{Int}(V)$.

There is a natural ‘almost-homeomorphism’ between 2^ω and $[0, 1]$: given an infinite sequence $x \in 2^\omega$ we have $0.x \in [0, 1]$ (interpreting the sequence as a decimal expansion in binary), and conversely given $y \in [0, 1]$ we can take the binary expansion of y containing infinitely many 0s, which gives us an element of 2^ω . Note that the problem of nonunique expansions only occurs for rationals, which are not 1-generic anyway. It is thus natural to ask if the notions of 1-genericity in these two spaces correspond via this mapping. The next proposition says this is indeed the case.

Proposition 2.5. For any irrational $x \in [0, 1]$ we have that x is 1-generic if and only if its (unique) binary expansion is 1-generic in 2^ω .

Proof. Let 2^ω_- be the set of infinite binary sequences which contain infinitely many 0s and infinitely many 1s. Then the ‘almost-homeomorphism’ given above in fact restricts to a homeomorphism to 2^ω_- and $[0, 1]_-$, where $[0, 1]_-$ is $[0, 1]$ without the dyadic rationals. Therefore, 1-genericity on 2^ω_- and $[0, 1]_-$ (which are defined as in Definition 2.4) coincide.

Note that 2^ω_- is dense in 2^ω and that $[0, 1]_-$ is dense in $[0, 1]$. Thus, it is enough if we can show that if $Y \subseteq X$ is such that Y is dense in X , then 1-genericity on X and Y coincide for elements $y \in Y$. Given a Π_1^0 class $V \subseteq X$, let $W = V \cap Y$. Then W is a Π_1^0 class in Y . Conversely, every Π_1^0 class $W \subseteq Y$ is of the form $W = V \cap Y$ by definition of the induced topology.

We claim that $\text{Int}_X(V) \cap Y = \text{Int}_Y(W)$. Clearly, $\text{Int}_X(V) \cap Y \subseteq \text{Int}_Y(W)$. Conversely, if we let $\text{Int}_Y(W) = U \cap Y$ for some open $U \subseteq X$, then $U \subseteq \text{Int}_X(V \cup (X \setminus Y))$. Towards a contradiction, assume that $U \cap (X \setminus V) \neq \emptyset$, then this is a nonempty open set. However, we also have $U \cap (X \setminus V) \subseteq X \setminus Y$, which contradicts the fact that Y is dense in X . Thus, we see that $U \subseteq V$, and therefore $U \subseteq \text{Int}_X(V)$. So, $\text{Int}_X(V) \cap Y = \text{Int}_Y(W)$.

So, we see that $y \notin V \setminus \text{Int}_X(V)$ if and only if $y \notin W \setminus \text{Int}_Y(W)$. This completes the proof. \square

3. EFFECTIVE BAIRE CLASS 1 FUNCTIONS

In this section we will discuss what it means for a function to be of effective Baire class 1, and discuss some of the basic properties of these functions. First, let us recall what it means for a function on the reals to be computable. Our definitions follow Moschovakis [9].

Definition 3.1. Let $f : [0, 1] \rightarrow \mathbb{R}$. We say that f is *computable* if for every basic open set U we have that $f^{-1}(U)$ is Σ_1^0 uniformly in U , i.e. if there exists a computable function $\alpha : \mathbb{Q} \times \mathbb{Q} \rightarrow \omega$ such that for all $q, r \in \mathbb{Q}$ we have that $f^{-1}((q, r))$ is equal to the Σ_1^0 class given by the index $\alpha(q, r)$.

Definition 3.1 is equivalent to the formulation with computable functionals, see e.g. the discussion in Pour-El and Richards [13].

Functions of effective Baire class 1 are obtained by weakening the above definition as follows.

Definition 3.2. A function $f : [0, 1] \rightarrow \mathbb{R}$ is of *effective Baire class 1* if for every basic open set U we have that $f^{-1}(U)$ is Σ_2^0 uniformly in U .

Replacing Σ_2^0 by Σ_2^0 in the above definition, we obtain what is known as a function of (*non-effective*) *Baire class 1*. Before we give an important example of an effective Baire class 1 function, let us first consider the following proposition, which gives an equivalent condition for a function to be of effective Baire class 1. This proposition mirrors the classical proposition which says that a function is of Baire class 1 if and only if it is a pointwise limit of continuous functions, see e.g. Kechris [7, p. 192]. (This does not hold for all Polish spaces; it holds for $f : X \rightarrow Y$ if either X is zero-dimensional or $Y = \mathbb{R}$.)

Proposition 3.3. *Let $f : [0, 1] \rightarrow \mathbb{R}$. The following are equivalent:*

- (i) f is of effective Baire class 1,
- (ii) f is the pointwise limit of a uniform sequence of computable functions, i.e. there exists a sequence f_0, f_1, \dots of functions from $[0, 1]$ to \mathbb{R} converging pointwise to f and a computable function $\alpha : \omega \times \mathbb{Q} \times \mathbb{Q} \rightarrow \omega$ such that for all $q, r \in \mathbb{Q}$ and all $n \in \omega$ we have that $f_n^{-1}((q, r))$ is equal to the Σ_1^0 class given by the index $\alpha(n, q, r)$.

Proof. (ii) \rightarrow (i): Let f_0, f_1, \dots be a sequence of uniformly computable functions converging to f and let U be any basic open set. Then $U = \bigcup_{i \in \omega, V_i \subseteq U} V_i$, where V_0, V_1, \dots is a computable enumeration of the closed intervals with rational endpoints. We claim:

$$f^{-1}(U) = \bigcup_{V_i \subseteq U} \bigcup_{n \in \omega} \bigcap_{m \geq n} f_m^{-1}(V_i),$$

which is clearly Σ_2^0 uniformly in U .

To prove the claim, let $x \in f^{-1}(U)$. Then $f(x) \in U$, so there exists $V_i \subseteq U$ such that $f(x) \in \text{Int}(V_i)$, say $(f(x) - \varepsilon, f(x) + \varepsilon) \subseteq V_i$. Let $n \in \omega$ be such that for every $m \geq n$ we have that $|f_m(x) - f(x)| < \varepsilon$. Then for every $m \geq n$ we have that $x \in f_m^{-1}(V_i)$, which proves the first inclusion.

Conversely, let $n \in \omega$, $V_i \subseteq U$ and $x \in \bigcap_{m \geq n} f_m^{-1}(V_i)$. Then for every $m \geq n$ we have $f_m(x) \in V_i$, and since V_i is closed we then also have $f(x) = \lim_{m \rightarrow \infty} f_m(x) \in V_i \subseteq U$, which completes the proof of the claim.

(i) \rightarrow (ii): This follows by effectivizing Kechris [7, Theorem 24.10]; this result is also mentioned (without proof) in Moschovakis [9, Exercise 3.E.14]. Since this implication is not used anywhere in this paper, we will not go into further detail. \square

Using this proposition, we can now give an important example of effective Baire class 1 functions: derivatives of computable functions. This also explains our interest in them.

Corollary 3.4. *Let $f : [0, 1] \rightarrow \mathbb{R}$ be a differentiable computable function. Then f' is of effective Baire class 1.*

Proof. Let $f_n(x) = 2^n(f(x + 2^{-n}) - f(x))$. To account for the problem that for $x + 2^{-n} > 1$ the value $f(x + 2^{-n})$ is not defined, we let $f(y) = -f(2 - y) + 2f(1)$ for $y > 1$ (i.e. we flip and mirror f on $[1, 2]$). Then the sequence f_0, f_1, \dots is uniformly computable and converges pointwise to f' , so f' is of effective Baire class 1 by Proposition 3.3. \square

4. CONTINUITY OF BAIRE CLASS 1 FUNCTIONS

At the basis of this section lies the following important classical result.

Theorem 4.1. (Baire) *Let $f : [0, 1] \rightarrow \mathbb{R}$ be of (*non-effective*) Baire class 1. Then the points of discontinuity of f form a meager Σ_2^0 set.*

Proof. See Kechris [7, Theorem 24.14] or Oxtoby [12, Theorem 7.3]. \square

We will now effectivize this result.

Theorem 4.2. *Let $f : [0, 1] \rightarrow \mathbb{R}$ be of effective Baire class 1. Then f is continuous at every 1-generic point.*

Proof. We effectivize the proof from Kechris [7, Theorem 24.14]. Let U_0, U_1, \dots be an effective enumeration of the basic open sets. Now f is continuous at x if and only if the inverse image of every neighborhood of $f(x)$ is a neighborhood of x . Thus, f is discontinuous at x if and only if there exists an open set U containing $f(x)$ such that every open set contained in $f^{-1}(U)$ does not contain x . Hence

$$\{x \in [0, 1] \mid f \text{ is discontinuous at } x\} = \bigcup_{n \in \omega} f^{-1}(U_n) \setminus \text{Int}(f^{-1}(U_n)).$$

Now, let x be such that f is discontinuous at x and let n be such that $x \in f^{-1}(U_n) \setminus \text{Int}(f^{-1}(U_n))$. Because f is of effective Baire class 1, we know that $f^{-1}(U_n)$ is Σ_2^0 . So, let $f^{-1}(U_n) = \bigcup_{i \in \omega} V_i$, where each V_i is Π_1^0 . Then it is directly verified that

$$f^{-1}(U_n) \setminus \text{Int}(f^{-1}(U_n)) \subseteq \bigcup_{i \in \omega} (V_i \setminus \text{Int}(V_i)).$$

Let i be such that $x \in V_i \setminus \text{Int}(V_i)$. Then x is not 1-generic by Definition 2.4. \square

Combining this result with the fact that derivatives of computable functions are of effective Baire class 1, we get the first implication of Theorem 1.1 as a consequence.

Theorem 4.3. *If $f : [0, 1] \rightarrow \mathbb{R}$ is a computable function, then f' is continuous at every 1-generic real.*

Proof. From Corollary 3.4 and Theorem 4.2. \square

5. FUNCTIONS DISCONTINUOUS AT NON-1-GENERIC

In this section we will prove the second implication of Theorem 1.1. To this end, we will build, for each Π_1^0 class V , a Volterra-style differentiable computable function whose derivative will fail to be continuous at the points whose non-1-genericity is witnessed by V . We have to be careful in order to make this function computable.

Theorem 5.1. *Let V be a Π_1^0 class. Then there exists a differentiable computable function $f : [0, 1] \rightarrow \mathbb{R}$ such that f' is discontinuous at every $x \in V \setminus \text{Int}(V)$.*

Proof. In the construction of f below, we first define auxiliary functions g and h .

Construction. We define an auxiliary function g , with the property that g is differentiable and computable, and g' is continuous on $(0, 1)$ and discontinuous at 0 and 1.

Define the function h on $[0, 1]$ by $h(0) = 0$ and

$$h(x) = x^2 \sin\left(\frac{1}{x^2}\right)$$

for $x > 0$. Then h is computable and differentiable, with derivative $h'(0) = 0$ and

$$h'(x) = 2x \sin\left(\frac{1}{x^2}\right) - 2\frac{1}{x} \cos\left(\frac{1}{x^2}\right)$$

when $x > 0$. Note that h' is discontinuous at $x = 0$. Fix a computable $x_0 \in (0, \frac{1}{2}]$ such that $h'(x_0) = 0$. Such an x_0 exists, because h' has isolated roots, and isolated

roots of computable functions are computable. Now define g on $[0, 1]$ by

$$g(x) = \begin{cases} 0 & \text{if } x = 0 \\ h(x) & \text{if } x \in (0, x_0] \\ h(x_0) & \text{if } x \in [x_0, 1 - x_0] \\ h(1 - x) & \text{if } x \in [1 - x_0, 1) \\ 0 & \text{if } x = 1. \end{cases}$$

Then g is a differentiable computable function, with derivative

$$g'(x) = \begin{cases} 0 & \text{if } x = 0 \\ h'(x) & \text{if } x \in (0, x_0] \\ 0 & \text{if } x \in [x_0, 1 - x_0] \\ -h'(1 - x) & \text{if } x \in [1 - x_0, 1) \\ 0 & \text{if } x = 1. \end{cases}$$

In particular, we see that g' is continuous exactly on $(0, 1)$. We will use g to construct f .

For the given Π_1^0 class V , let $U = [0, 1] \setminus V$, and fix computable enumerations q_0, q_1, \dots and r_0, r_1, \dots of rational numbers in $[0, 1]$ such that $U = \bigcup_{n \in \omega} [q_n, r_n]$ and such that the (q_n, r_n) are pairwise disjoint. We will construct f as a sum of a sequence f_0, f_1, \dots of uniformly computable functions. We define f_n by:

$$(5.1) \quad f_n(x) = \begin{cases} 0 & \text{if } x \in [0, q_n] \\ \frac{r_n - q_n}{2^n} g\left(\frac{x - q_n}{r_n - q_n}\right) & \text{if } x \in [q_n, r_n] \\ 0 & \text{if } x \in [r_n, 1]. \end{cases}$$

Finally, we let $f = \sum_{n=0}^{\infty} f_n$.

Verification. We first show that f is computable. To this end, first observe that each f_n is supported on (q_n, r_n) , and therefore the supports of the different f_n are disjoint. Furthermore, each f_n is bounded by 2^{-n} .

Let (a, b) be a basic open subset of \mathbb{R} . We distinguish two cases. First, assume $0 \notin (a, b)$. We assume $a > 0$, the case $b < 0$ is proven in a similar way. Let $n \in \omega$ be such that $2^{-n} < a$. Then, since the supports of the f_m are disjoint, and each f_m is bounded by 2^{-m} , we have

$$f^{-1}((a, b)) = (f_0 + \dots + f_n)^{-1}((a, b)),$$

which is Σ_1^0 because a finite sum of computable functions is computable.

In the second case, we have $0 \in (a, b)$. Let n be such that $|a|, |b| \geq 2^{-n}$. Then, again because the supports of the f_m are disjoint, we see that if x is not in the support of any f_m for $m \leq n$ then certainly $f(x) \in (a, b)$. Therefore we have

$$f^{-1}((a, b)) = (f_0 + \dots + f_n)^{-1}((a, b)) \cup \bigcap_{m \leq n} ([0, 1] \setminus [q_m, r_m]),$$

which is also Σ_1^0 . It is clear that the case distinction is uniformly computable, so it follows that f is computable.

Next, we check that f is differentiable. We first note that every f_n is differentiable, because g is differentiable. Let $x \in [0, 1]$. We distinguish two cases. First, if x is in some (q_n, r_n) then it is immediate that f is differentiable at x with derivative $f'_n(x)$, because the intervals (q_n, r_n) are disjoint. Next, we consider the case where x is not

in any interval (q_n, r_n) . Note that in this case we have $f(x) = 0$. Fix $m \in \omega$. Then we have:

$$\lim_{y \rightarrow x} \left| \frac{f(y)}{y-x} \right| \leq \lim_{y \rightarrow x} \left| \frac{(f_0 + \dots + f_m)(y)}{y-x} \right| + \lim_{y \rightarrow x} \left| \frac{(f_{m+1} + f_{m+2} + \dots)(y)}{y-x} \right|.$$

Because $f_0 + \dots + f_m$ is differentiable at x with derivative 0, this is equal to:

$$(5.2) \quad \lim_{y \rightarrow x} \left| \frac{(f_{m+1} + f_{m+2} + \dots)(y)}{y-x} \right|.$$

To show that this limit is 0, we will prove that it is bounded by $\frac{1}{2^m(1-x_0)}$ for every m . Let $y \in [0, 1]$ be distinct from x . Let us assume that $x < y$; the other case is proven in the same way. If y is not in any (q_n, r_n) for $n \geq m+1$ then $(f_{m+1} + f_{m+2} + \dots)(y) = 0$. Otherwise, there is exactly one such n . Then:

$$\left| \frac{(f_{m+1} + f_{m+2} + \dots)(y)}{y-x} \right| = \left| \frac{f_n(y)}{y-x} \right| \leq \left| \frac{f_n(y)}{y-q_n} \right|,$$

where the last inequality follows from the fact that x does not lie in (q_n, r_n) . We distinguish three cases. First, if $z = \frac{y-q_n}{r_n-q_n} \in (0, x_0]$, then

$$\left| \frac{f_n(y)}{y-q_n} \right| = \left| \frac{2^{-n}(r_n-q_n)g(z)}{y-q_n} \right| = |2^{-n}z \sin(z^{-2})| \leq 2^{-n} \leq \frac{1}{2^m(1-x_0)}.$$

Next, if $z \in [x_0, 1-x_0]$ (which is nonempty because $x_0 \leq \frac{1}{2}$), then

$$\left| \frac{f_n(y)}{y-q_n} \right| \leq \frac{2^{-n}(r_n-q_n)x_0^2}{y-q_n} = \frac{2^{-n}x_0^2}{z} \leq x_0 2^{-n} \leq \frac{1}{2^m(1-x_0)}$$

where we use the fact that $z \geq x_0$. Finally, if $z \in [1-x_0, 1]$, then

$$\left| \frac{f_n(y)}{y-q_n} \right| = \left| \frac{2^{-n}(r_n-q_n)h(1-z)}{y-q_n} \right| \leq \frac{1}{2^n z} \leq \frac{1}{2^n(1-x_0)} \leq \frac{1}{2^m(1-x_0)}.$$

Combining this with (5.2) we see that $\lim_{y \rightarrow x} \left| \frac{f(y)}{y-x} \right| \leq \frac{1}{2^m(1-x_0)}$. Since m was arbitrary this shows that f is differentiable at x , with derivative $f'(x) = 0$.

Finally, we need to verify that f' is discontinuous at x for all $x \in V \setminus \text{Int}(V)$. Therefore, let $x \in V \setminus \text{Int}(V)$. Then every open set W containing x has nonempty intersection $W \cap U$ (recall that $U = [0, 1] \setminus V$), but this intersection does not contain x . We have shown above that $f'(x) = 0$. We will show that for every open interval I containing x there is a point $y \in I$ such that $f'(y) \leq -1$, which clearly shows that f' cannot be continuous at x . Fix an open interval I containing x . Then $I \cap U \neq \emptyset$, so there is an $n \in \omega$ such that $I \cap [q_n, r_n]$ is nonempty. Note that I contains x and therefore I cannot be a subinterval of $[q_n, r_n]$. Therefore there exists a $q_n < s < r_n$ such that either $[q_n, s] \subseteq I$ or $(s, r_n] \subseteq I$. We will assume the first case; the second case is proven in a similar way.

Note that on $[q_n, s)$ the function f' is equal to f'_n . For $y \in (q_n, s)$ we thus have:

$$f'(y) = 2^{-n}g'((y-q_n)/(r_n-q_n)).$$

So, we need to show that there is a $y \in (q_n, s)$ such that $g'((y-q_n)/(r_n-q_n)) \leq -2^n$, or equivalently, that there is a $z \in (0, (s-q_n)/(r_n-q_n))$ such that $g'(z) \leq -2^n$. Without loss of generality, $(s-q_n)/(r_n-q_n) < x_0$. Let $k \geq n$ be such that $2^{-k} \leq \frac{s-q_n}{r_n-q_n}$. Then:

$$\begin{aligned} g'(1/(2^k \sqrt{\pi})) &= \frac{1}{2^{k-1} \sqrt{\pi}} \sin(2^{2k} \pi) - 2^{k+1} \sqrt{\pi} \cos(2^{2k} \pi) \\ &= -2^{k+1} \sqrt{\pi} \leq -2^k \leq -2^n. \end{aligned}$$

This completes the verification. \square

Theorem 5.2. *If $x \in [0, 1]$ is such that every differentiable computable function $f : [0, 1] \rightarrow \mathbb{R}$ has continuous derivative at x , then x is 1-generic.*

Proof. If x is not 1-generic, then there is a Π_1^0 class V such that $x \in V \setminus \text{Int}(V)$. Applying Theorem 5.1 to V gives a differentiable computable function f for which f' is discontinuous at x . \square

6. n -GENERICITY

The notion of 1-genericity (Definition 2.1) corresponds to the first level of the arithmetical hierarchy. Higher genericity notions can be defined using forcing conditions from higher levels of the arithmetical hierarchy. As for 1-genericity, an equivalent formulation can be given as follows, see Jockusch [6]:

Definition 6.1. An element $x \in 2^\omega$ is n -generic if x meets every Σ_n^0 set of strings $A \subseteq 2^{<\omega}$ that is dense along x .

As usual, let \emptyset' denote the halting set, and let $\emptyset^{(n)}$ denote the n -th jump. Since a Σ_n^0 set of strings is the same as a Σ_1^0 set of strings relative to $\emptyset^{(n-1)}$, a set is n -generic if and only if it is 1-generic relative to $\emptyset^{(n-1)}$.

Corollary 2.3 relativizes to:

Proposition 6.2. *For any $x \in 2^\omega$ we have that x is n -generic if and only if for every $\Pi_1^{0, \emptyset^{(n-1)}}$ class $V \subseteq 2^\omega$ we have $x \notin V \setminus \text{Int}(V)$.*

Note that in general a $\Pi_1^{0, \emptyset^{(n-1)}}$ class in 2^ω is not the same as a Π_n^0 class, since the latter need not even be closed. (And even if one *assumes* that the class is closed the notions are not the same, see [5, p76].)

Given this equivalence, we can now generalize Definition 2.4 to:

Definition 6.3. Let $x \in [0, 1]$. We say that x is n -generic if for every $\Pi_1^{0, \emptyset^{(n-1)}}$ class $V \subseteq [0, 1]$ we have $x \notin V \setminus \text{Int}(V)$.

Further justification for this definition comes from the fact that Proposition 2.5 relativizes: An irrational $x \in [0, 1]$ is n -generic according to Definition 6.3 if and only if its binary expansion is n -generic in 2^ω .

It is straightforward to check that the results of all the previous sections relativize to an arbitrary oracle A . This gives the following relativized version of Theorem 1.1:

Theorem 6.4. *A real $x \in [0, 1]$ is 1-generic relative to A if and only if for every differentiable A -computable function $f : [0, 1] \rightarrow \mathbb{R}$, f' is continuous at x .*

Taking $A = \emptyset^{(n-1)}$, this immediately gives the following characterization of n -genericity:

Corollary 6.5. *A real $x \in [0, 1]$ is n -generic if and only if for every differentiable $\emptyset^{(n-1)}$ -computable function $f : [0, 1] \rightarrow \mathbb{R}$, f' is continuous at x .*

Also, taking all n together, we see that a real x is arithmetically generic if and only if every differentiable arithmetical function has continuous derivative at x .

7. MULTIPLY DIFFERENTIABLE FUNCTIONS

We have characterized 1-genericity using the continuity of the derivatives of (once) differentiable computable functions. One might wonder: what kind of effective genericity for x corresponds to every twice differentiable, computable function having continuous second derivative at x ? Or, more generally, what corresponds to every n times differentiable, computable function having continuous n th derivative at x ?

It turns out that the answer is always 1-genericity. To show this we will need the following proposition, which essentially tells us that the case for $n > 2$ collapses to the case $n = 2$.

Proposition 7.1. *Let $f : [0, 1] \rightarrow \mathbb{R}$ be computable and twice continuously differentiable. Then f' is computable.*

Proof. See e.g. Pour-El and Richards [13, Theorem 1.2]. \square

If the second derivative of a computable function exists, it is easy to see that it is of effective Baire class 2 (i.e. a pointwise limit of a computable sequence of functions of effective Baire class 1), by similar arguments as in the proof of Corollary 3.4. However, using the following proposition we can easily see that the second derivative of a computable function is in fact of effective Baire class 1.

Proposition 7.2. *Let $f : [0, 1] \rightarrow \mathbb{R}$ be twice differentiable. Then*

$$f''(x) = \lim_{h \rightarrow 0} \frac{f(x+h) + f(x-h) - 2f(x)}{h^2}.$$

Proof. See e.g. Rudin [14, p. 115]. \square

Theorem 7.3. *Fix $n \geq 1$. Then a real $x \in [0, 1]$ is 1-generic if and only if every n times differentiable, computable function $f : [0, 1] \rightarrow \mathbb{R}$ has continuous n th derivative at x .*

Proof. For $n = 1$ this is exactly Theorem 1.1. So, we may assume $n \geq 2$. First, if $x \in [0, 1]$ is not 1-generic, then by Theorem 1.1 there is a differentiable, computable function $g : [0, 1] \rightarrow \mathbb{R}$ such that g' is not continuous at x . Now let $h_1 = g$ and let h_i be a computable antiderivative of h_{i-1} for $2 \leq i \leq n$ (which exists by Ko [8, Theorem 5.29]). Then, if we let $f = h_n$, we see that f is an n times differentiable, computable function such that $f^{(n)}$ is discontinuous at x .

Conversely, if f is an n times differentiable, computable function, then $f^{(n-2)}$ is computable by Proposition 7.1. So, $f^{(n)}$ is of effective Baire class 1 by Proposition 7.2. Thus, if $f^{(n)}$ is discontinuous at x , then x is not 1-generic by Theorem 4.2. \square

8. COMPLEXITY THEORETIC CONSIDERATIONS

In this section we discuss polynomial time computable real functions. The theory of these functions is developed in Ko [8], to which we refer the reader for the basic results and definitions. Briefly, a function $f : [0, 1] \rightarrow \mathbb{R}$ is polynomial time computable if for any $x \in [0, 1]$ we can compute an approximate value of $f(x)$ to within an error of 2^{-n} in time n^k for some constant k .

Most of the common functions from analysis, such as rational functions and the trigonometric functions, as well as their inverses, are all polynomial time computable, see e.g. Brent [2] and Weihrauch [15]. Also, the polynomial time computable functions are closed under composition. With this knowledge, it is not difficult to see that the construction of the function f in section 5 can be modified to yield a polynomial time computable function, rather than just a computable one. For this it is also needed that the complement of the Π_1^0 class V from Theorem 5.1 can be represented by a polynomial time computable set of strings. This is similar to the fact that every nonempty computably enumerable set is the range of a polynomial time computable function, simply by sufficiently slowing down the enumeration. Since the enumeration of $U = [0, 1] \setminus V$ in the proof of Theorem 5.1 is now slower, the definition of f_n in (5.1) has to be adapted by replacing 2^n by $2^{t(n)}$, where $t(n)$ is the stage at which the interval (q_n, r_n) is enumerated into U . This modification ensures that the functions f_n are uniformly polynomial time computable, so that

also the function $f = \sum_{n=0}^{\infty} f_n$, is polynomial time computable. Thus we obtain the following strengthening of Theorem 5.1:

Theorem 8.1. *Let V be a Π_1^0 class. Then there exists a differentiable polynomial time computable function $f : [0, 1] \rightarrow \mathbb{R}$ such that f' is discontinuous at every $x \in V \setminus \text{Int}(V)$.*

We now have the following variant of Theorem 1.1:

Theorem 8.2. *A real $x \in [0, 1]$ is 1-generic if and only if for every differentiable polynomial time computable function $f : [0, 1] \rightarrow \mathbb{R}$, f' is continuous at x .*

Proof. The “only if” direction is immediate from Theorem 1.1. For the “if” direction; if x is not 1-generic, then there is a Π_1^0 class V such that $x \in V \setminus \text{Int}(V)$. Theorem 8.1 then gives a differentiable polynomial time computable function f for which f' is discontinuous at x . \square

REFERENCES

- [1] V. Brattka, J. S. Miller, and A. Nies, *Randomness and differentiability* arXiv:1104.4465 [math.LO], 2011.
- [2] R. P. Brent, *Fast multiple-precision evaluation of elementary functions*, Journal of the ACM 23(2) (1976) 242–251.
- [3] A. M. Bruckner and J. L. Leonard, *Derivatives*, The American Mathematical Monthly 73(4) Part 2: Papers in Analysis (1966) 24–56.
- [4] O. Demuth, *The differentiability of constructive functions of weakly bounded variation on pseudo numbers*, Commentationes Mathematicae Universitatis Carolinae 16(3)(1975) 583–599. (In Russian.)
- [5] R. G. Downey and D. R. Hirschfeldt, *Algorithmic randomness and complexity*, Springer, 2010.
- [6] C. G. Jockusch, *Degrees of generic sets*, in: Recursion Theory: its Generalisations and Applications, edited by F. R. Drake and S. S. Wainer, Cambridge University Press, 1980, 110–139.
- [7] A. S. Kechris, *Classical descriptive set theory*, Springer-Verlag, 1995.
- [8] K. Ko, *Complexity theory of real functions*, Birkhäuser, 1991.
- [9] Y. N. Moschovakis, *Descriptive set theory*, second edition, Mathematical Surveys and Monographs, Vol. 155, American Mathematical Society, 2009.
- [10] P. G. Odifreddi, *Classical recursion theory*, Vol. 1, Studies in Logic and the Foundations of Mathematics Vol. 125, North-Holland, 1989.
- [11] P. G. Odifreddi, *Classical recursion theory*, Vol. 2, Studies in Logic and the Foundations of Mathematics Vol. 143, North-Holland, 1999.
- [12] J. C. Oxtoby, *Measure and category*, second ed., Springer-Verlag, 1980.
- [13] M. B. Pour-El and J. I. Richards, *Computability in Analysis and Physics*, Perspectives in Mathematical Logic, Vol. 1, Springer-Verlag, 1989.
- [14] W. Rudin, *Principles of Mathematical Analysis*, third ed., McGraw-Hill Book Company, 1976.
- [15] K. Weihrauch, *Computable analysis, an introduction*, Springer-Verlag, 2000.

(Rutger Kuiper) RADBOD UNIVERSITY NIJMEGEN, DEPARTMENT OF MATHEMATICS, P.O. BOX 9010, 6500 GL NIJMEGEN, THE NETHERLANDS.

E-mail address: `r.kuiper@math.ru.nl`

(Sebastiaan A. Terwijn) RADBOD UNIVERSITY NIJMEGEN, DEPARTMENT OF MATHEMATICS, P.O. BOX 9010, 6500 GL NIJMEGEN, THE NETHERLANDS.

E-mail address: `terwijn@math.ru.nl`